

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
29 November 2001 (29.11.2001)

PCT

(10) International Publication Number
WO 01/89586 A2

- (51) International Patent Classification⁷: **A61K 51/04** 63017 (US). **DYSZLEWSKI, Mary, E.** [US/US]; 6 West Walling Drive, Creve Coeur, MO 63141 (US). **WEBB, Elizabeth, G.** [US/US]; 220 Shelburne Drive, St. Charles, MO 63301 (US).
- (21) International Application Number: PCT/US01/15670
- (22) International Filing Date: 8 May 2001 (08.05.2001) (74) Agent: **BOONE, Jeffrey, S.**; 675 McDonnell Boulevard, P.O. Box 5840, St. Louis, MO 63134 (US).
- (25) Filing Language: English
- (26) Publication Language: English (81) Designated States (*national*): JP, US.
- (30) Priority Data: 09/576,960 24 May 2000 (24.05.2000) US (84) Designated States (*regional*): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR).
- (71) Applicant (*for all designated States except US*): **MALLINCKRODT INC.** [US/US]; 675 McDonnell Boulevard, P.O. Box 5840, St. Louis, MO 63134 (US). Published:
— *without international search report and to be republished upon receipt of that report*
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **PIPES, David, W.** [US/US]; 14517 Debbelham Lane, Chesterfield, MO
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

BEST AVAILABLE COPY

WO 01/89586 A2

(54) Title: FORMULATION OF TC AND RE CARBONYL COMPLEXES USING STANNOUS ION AS THE REDUCTANT FOR PERTECHNETATE AND PERRHENATE

(57) Abstract: The invention relates to novel aminocarboxylate ligands that are suitable for complexing with a radionuclide, and are useful as therapeutic agents and as imaging agents for diagnostic purposes. In accordance with the present invention, a method of preparing a compound of formula $fac-[M(CO)_3(OH_2)_3]^+$ (I) wherein M is Mn, ^{99m}Tc , ^{186}Re or ^{188}Re , involves reacting a metal in permetallate form with carbon monoxide and stannous ion. The compound of formula (I) can be reacted with a ligand L_x , to form a compound of the formula $fac-[M(CO)_3L_x]^n$ (II) wherein M is as defined above, L_x is a monodentate or multidentate ligand or a mixture of these ligands, and n is a charge of the ligand L_x increased with one + charge. The invention also is directed to novel compounds, and kits for carrying out the disclosed methods.

TITLE OF THE INVENTION

FORMULATION OF Tc AND Re CARBONYL COMPLEXES USING STANNOUS ION AS THE REDUCTANT FOR PERTECHNETATE AND PERRHENATE

5 FIELD OF THE INVENTION

The present invention relates to a novel method for preparing Mn, Tc and Re carbonyl complexes which are useful such as for imaging and therapeutic agents.

BACKGROUND OF THE INVENTION

10 Scintigraphic imaging and similar radiographic techniques for visualizing tissues *in vivo* are finding ever-increasing application in biological and medical research and in diagnostic and therapeutic procedures. Generally, scintigraphic procedures involve the preparation of radioactive agents which, upon introduction to a biological subject, become localized in the specific organ, tissue or skeletal structure of choice. When so localized,
15 traces, plots or scintiphotos depicting the *in vivo* distribution of radiographic material can be made by various radiation detectors, e.g., traversing scanners and scintillation cameras. The distribution and corresponding relative intensity of the detected radioactive material not only indicates the space occupied by the targeted tissue, but also indicates a presence of receptors, antigens, aberrations, pathological conditions, and the like.

20 Technetium-99m (^{99m}Tc) is a radionuclide which is widely known for its uses in tissue imaging agents. Due to its safety and ideal imaging properties, this radionuclide is conveniently available commercially in the oxidized pertechnetate form ($^{99m}\text{TcO}_4^-$) hereinafter "pertechnetate-Tc99m". However, pertechnetate will not complex with the most commonly used biological carriers for radionuclide tissue imaging. Thus, technetium-labeled
25 imaging agents are generally prepared by admixing a pertechnetate-Tc99m isotonic saline solution, a technetium reductant (reducing agent) such as stannous chloride or sodium dithionite, and a chelate conjugated to the desired peptide carrier agent for targeting the organ of interest. Alternatively, an intermediate transfer liquid-technetium 99m complex may be prepared prior to addition to the chelate-biological molecule to maintain the oxidation state
30 within a desired level. Examples of such include ^{99m}Tc -tartrate or ^{99m}Tc -gluconate.

Another problem is that technetium-containing scintigraphic imaging agents are known to be unstable in the presence of oxygen, primarily since oxidation of the reductant and/or the technetium-99m destroys the reduced technetium-99m/targeting carrier complex. Accordingly, such imaging agents are generally made oxygen-free by saturating the


compositions with oxygen-free nitrogen gas or by preparing the agents in an oxygen-free atmosphere. Stabilization of imaging agents can also be achieved through chemical means. U.S. Pat. No. 4,232,000, Fawzi, issued Nov. 4, 1980, discloses the use of gentisyl alcohol as a stabilizer for technetium imaging agents. Similarly, U.S. Pat. No. 4,233,284, Fawzi, issued Nov. 11, 1980, discloses the use of gentisic acid as a stabilizer.

In published PCT Application No. PCT/US98/07979 (International Publication No. WO 98/48848), which is incorporated herein in its entirety by reference, a method was disclosed for preparing a compound of the general formula (I): $fac-[M(CO)_3(OH_2)_3]^+$ wherein M is Mn, ^{99m}Tc , ^{186}Re or ^{188}Re , by reacting a metal in the permethylate form with carbon monoxide and a reducing agent, characterized in that a mixture of a base, a reducing agent soluble in water but not substantially decomposed by water, and optionally a stabilizing agent is dissolved in a water containing solvent system containing a solution of the metal in the permanganate, pertechnetate or perrhenate form in the presence of carbon monoxide and optionally in the presence of a halide. The ligands disclosed for labeling biologically active molecules have a tendency to stabilize metals in their low oxidation states. These ligands have in common the presence of low-lying vacant orbitals of the correct symmetry to form pi-bonds by accepting electrons from filled metal d-orbitals, a phenomenon known as backbonding. The ligands indicated in the patent application include isonitriles, phosphines, thioethers, Schiff bases, and pyridine-, imidazole-, and pyrazole-type groups. In particular, the amino acid histidine is indicated as an ideal chelate. For some purposes a problem with using histidine and other unsaturated organic molecules as chelates is that the resulting labeled compound is highly lipophilic resulting in high liver and blood uptake. The predominant hepatobiliary uptake and clearance are for some purposes undesirable characteristics for the targeted imaging agents.

The publications and other materials used herein to illuminate the background of the invention or provide additional details respecting the practice, are incorporated by reference, and for convenience are respectively grouped in the appended List of References.

SUMMARY OF THE INVENTION

In accordance with the present invention, a method of preparing a compound of formula



3

wherein M is Mn, ^{99m}Tc , ^{186}Re or ^{188}Re ,

involves reacting a metal in permethylate form with carbon monoxide and a reducing agent, wherein the reducing agent comprises a stannous ion. The compound of formula (I) can be reacted with a ligand L_x to form a compound of the formula



wherein M is as defined above, L_x is a monodentate or multidentate ligand or a mixture of these ligands, and n is a charge of the ligand L_x increased with one + charge. The invention also is directed to kits for performing the disclosed methods.

10 DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to an improved method for synthesizing Tc(I) complexes using stannous ion as the reducing agent. For formulating Tc(I) complexes, e.g., $[\text{Tc}(\text{CO})_3(\text{OH}_2)_3]^+$, the prior art methods have successfully used only the powerful reductants of borohydrides (BH_3 or BH_4^-). Attempts to use other reductants including stannous ion
 15 (Sn^{2+}) have resulted in labeling yields of approximately only 50%. The use of the more common Sn^{2+} in radiopharmaceutical kits has certain advantages such as a wide pH range of use, known toxicity, familiarity with the FDA and perhaps better adaptability between Tc and Re formulations. In addition, stannous ion is less likely to interfere with the biological substrate or ligands which are ultimately labeled. The instant invention discloses a method of
 20 using Sn^{2+} for the preparation of Tc-carbonyl complexes with yields of >80%, a clear improvement of the previous work.


It has normally been felt that to form Tc in oxidation state I as is the case of the Tc-carbonyl complexes formed according to this disclosure, a very strong reductant is needed, one with a negative value in the electrochemical series. The starting Tc compound is
 25 pertechnetate, TcO_4^- , which has Tc in oxidation state VII. To form the desired Tc-carbonyl complexes, it is necessary to reduce the Tc(VII) down to Tc(I).

Stannous ion is normally not thought of as a strong reductant because its electrochemical half equation is not very negative. In the presence of a 0.1 N acid, the electrochemical equation is:



The effect of pH upon this equation is usually forgotten. At a pH (>7) the reducing strength of stannous ion is changed as a result of a change in the reductant's

mechanism. In a basic solution, a stannous hydroxide chemical species is present with a different mechanism and reductant strength which is capable of reducing the Tc(VII) down to Tc(I), especially with the aid of an appropriate transfer ligand such as pyrophosphate or another ligand such as tartrate, gluceptate, methylenediphosphonate or hydroxyethyldiphosphonate. In a basic solution (0.1 N NaOH, pH=13), the half equation is:



This shows that stannous ion is a strong reductant at a basic pH. This is comparable to the electrochemical half equation for borohydride which is the reductant used in the prior art to form the desired Tc carbonyl compounds. Borohydride has an EE = -1.2 Volts at pH 13.

Methods are disclosed for preparation of facial metal tricarbonyl compounds and further coordinated facial metal tricarbonyl compounds. The invention further relates to the use of said facial metal tricarbonyl compounds in the labeling of biologically active substrates and other ligands, and to a kit for preparing a facial metal tricarbonyl compound or further co-ordinated facial metal tricarbonyl compounds.

It is known in the art (Alberto et al., 1994a) that facial metal tricarbonyl complexes of radioactive metals of group 7B of the periodic table are very convenient starting materials for substitution reactions in organic solvents as well as in water, as these compounds are stable in water for weeks, even if exposed to air. Therefore said compounds will be very useful for the labeling of biologically active substrates, such as amino acids, peptides, proteins, sugars and any receptor binding molecules. A major drawback, however, of these compounds until now is that they have only been available from high temperature carbonylation reactions and with the aid of the pyrophoric and toxic and therefore dangerous reducing agent BH_3 (Alberto et al., 1994b).

It is the objective of the present invention to provide for a method of preparing facial metal tricarbonyl compounds of (radioactive) metals of group 7B with the aid of easily available and low-toxic starting materials at moderate temperature and at normal pressure of CO, in a reasonable time and with high yield.

Such a method will be a powerful tool that can be used for the synthesis of diagnostic and therapeutic agents, especially for the synthesis of said diagnostic and therapeutic agents derived from radioactive metals with a short lifetime, in order to have access to these labeled compounds in poorly equipped hospital laboratories. When the above mentioned diagnostic agent is labeled with a radionuclide it can be detected by the so-called single photon emission

computerized tomography (SPECT), when it is labeled with a paramagnetic metal atom it can be detected by magnetic resonance imaging.

The above-defined objective can be achieved, according to the present invention, by a method of preparing a compound of the general formula



wherein M is Mn, ^{99m}Tc , ^{186}Re or ^{188}Re ,

by reacting a metal in the permethylate form (MO_4^- form) with carbon monoxide and a reducing agent comprising Sn^{2+} .

The metal M is preferably ^{99m}Tc , ^{186}Re or ^{188}Re , as these radionuclides, when used in diagnostic or therapeutic agents, have the advantage that they can be applied in very low concentrations, which minimizes the risk of toxicity.

It is very surprising that a quantitative reduction of permethylates in water containing solvent systems can be achieved at moderate temperature and in reasonable times with reducing agents that are nucleophilic and that are generally considered as less reactive than the electrophilic reducing agent BH_3 known in the art.

The method of the invention can be easily performed just by mixing the permethylate solution with the other reagents in the presence of carbon monoxide. The permethylate solution may optionally contain halide ions needed for the elution of the permethylate from a generator. The carbon monoxide may be supplied by using a closed system with an atmosphere containing a sufficient amount of carbon monoxide, or by flushing the carbon monoxide gas through the solution. Preferably the gas is substantially pure carbon monoxide. For the reduction the reducing agent is one comprising stannous ions. In preferred embodiments, the mixture includes lactose which is present as a bulking agent.

The reducing agent is reacted with the permethylate in a molar ratio higher than 3, preferably higher than 10 and most preferably higher than 100. The reduction reaction can be performed at a temperature between 20EC and 150EC. The preferred reaction temperature approximately 100EC. The heating of the reaction mixture can be performed in the normal way but also by microwave heating. The reaction can also be performed by the application of ultrasound, e.g., by carrying out the reactions in an ultrasonic bath at a temperature, normally leading to the same reaction rate at lower reaction temperature.

The compound of the general formula (I) obtained is very suitable for the labeling of biologically active substrates, such as amino acids, peptides, proteins, sugars, small receptor binding molecules or cells.

Examples of peptides that may be labeled are growth factors, somatostatin, bombesin, insulin, LHRH, gastrin, gastrin releasing peptide, thyrotropin releasing hormone, thyroid stimulating hormone, prolactin, vasoactive intestinal peptide (VIP), pituitary adenylate cyclase-activating polypeptide (PACAP), angiotensin, neurotensin, interferons, IL-1, IL-4 and IL-6, monoclonal antibodies and their analogues and derivatives. After labeling with a suitable labeling substance these peptides can, e.g., be used in the detection and localization of treatment of malignant human tumors.


Examples of sugars that may be labeled are glucose and deoxyglucose and derivatives of said compounds.

Small receptor binding molecules are defined as non-peptide molecules which are binding to a receptor and normally have a molecular mass below approximately 500 Daltons.

Examples of small receptor binding molecules that may be labeled are substances for the serotonergic system as described in WO 96/30054, or substances for the dopaminergic system (e.g., raclopride, 3-CIT, lisuride), for the cholinergic system (e.g., epibatidine), for the glutaminergic system (e.g., memantine) or for the benzodiazepine system (e.g., flumazenil, iomazenil). Examples of metabolic active molecules that may be labeled are DOPA, Tyrosine, mIBG, MAO-I and analogues thereof.

Examples of cells that may be labeled are red and white blood cells.

As a result of the labeling of (biologically active) substrates with a compound of the general formula I, a further coordinated compound of the general formula



is obtained wherein:

M is Mn, ^{99m}Tc, ¹⁸⁶Re or ¹⁸⁸Re;
L₁ is a monodentate ligand;
L₂ is a bidentate ligand;

L_3 is a tridentate ligand; and

n is the sum of the charge of the ligands L_1 , L_2 and L_3 increased with one + charge.

When the ligand L_1 , L_2 or L_3 before and/or after labeling with the facial metal tricarbonyl compound is the biologically active molecule, the present invention gives easy access to compounds that directly can be used as a diagnostic and therapeutic agent.

Examples of monodentate ligands within the definition of L_1 , L_2 and L_3 are (biologically active) substrates bearing groups such as phosphines, isonitriles, nitriles, imidazoles, thioethers and pyridine-like aromatic amines.

Examples of bidentate ligands within the definition of L_2 and L_3 are (biologically active) substrates bearing pyridine, imidazole or pyrazole groups, such as histidine, histamine, functionalized imidazole systems, bidentate thioethers, bidentate isocyanides, Schiff-base type ligands and picolinic acid.

Examples of tridentate ligands within the definition of L_3 are tris-pyrazolyl borate, tris-pyrazolylmethane, tris-imidazolyl borate, tris-pyrazolylmethane, 1,4,7-trithiacyclononane (9-aneS₃) and triazacyclononane (9-aneN₃), histidine, methionine, cysteine derivatized at the thiol group to give a thioether and cyclopentadienyl derivatives.

In some cases it may be advantageous to prepare the radiolabelled bioactive compound in one step. This objective can be achieved according to the present invention, with a method of preparing a compound of the general formula



wherein:

M is Mn, ^{99m}Tc, ¹⁸⁶Re or ¹⁸⁸Re;

L_1 is a monodentate ligand;

L_2 is a bidentate ligand;

L_3 is a tridentate ligand; and

n is the sum of the charge of the ligands L_1 , L_2 and L_3 increased with one + charge;

characterized in that a mixture of a base, ligands L_1 or L_2 or L_3 , a stannous ion and optionally a stabilizing agent is dissolved in a water containing solvent system containing a solution of

the metal in the permanganate, pertechnetate or perrhenate form in the presence of carbon monoxide and optionally in the presence of halide.

Especially in the case of radiolabelled compounds it is frequently impossible to put the ready-for-use composition at the disposal of the user, in connection with the often poor shelf life of the radiolabelled compound and/or the short half-life of the radionuclide used. In such cases the user will carry out the labeling reaction with the metal in the clinical hospital or laboratory. For this purpose the various reaction ingredients are then offered to the user in the form of a so-called "kit". It will be obvious that the manipulations necessary to perform the desired reaction should be as simple as possible to enable the user to prepare from the kit the radioactive labeled composition by using the facilities that are at his disposal. Therefore the invention also relates to a kit for preparing a labeling composition, which labeling composition contains a compound of formula I as the labeling agent.

Such a kit for the labeling of a biologically active substrate, according to the present invention, comprises (i) a reducing agent comprising stannous ion, (ii) if desired, a stabilizing agent and/or a chelator, e.g., pyrophosphate or glucoheptonate, and (iii) if desired one or more inert pharmaceutically acceptable carriers and/or formulating agents and/or adjuvants, e.g., lactose or inositol, at least one of said ingredients (i) to (iii) being stored in a container having an atmosphere containing a sufficient amount of carbon monoxide, or said ingredients (i) to (iii) being independently combined, and (iv) instructions for use with a prescription for reacting the ingredients of the kit with a metal selected from the group consisting of Mn, ^{99m}Tc , ^{186}Re or ^{188}Re in the form of permetallate solution. Preferably, the kit comprises a lyophilized formulation including pyrophosphate or gluceptate, SnCl_2 , and lactose, the mixture being sealed in a container having a headspace comprising carbon monoxide, most preferably substantially pure carbon monoxide. In other embodiments, the kit can include a metal (M) as defined above. In still further embodiments, the kit can include a ligand (L_x), which preferably is a multidentate aminopolycarboxylate ligand.

It is the merit of the present invention, disclosing an easy way of preparing facial tricarbonyl metal compounds within a time-frame that is reasonable compared with the half-life time of the radioactive isotopes involved, and with high yields, that a kit can be prepared for the labeling of biologically active substrates with said facial tricarbonyl metal compounds.

In some cases it may be advantageous to enclose a bioactive substrate in the kit so that a kit is obtained for the preparation of a radiopharmaceutical composition.

Alternatively the biologically active compound is formed upon the reaction of the ligand with the facial metal tricarbonyl compound.

5 Such a kit for the preparation of a diagnostic and therapeutic pharmaceutical composition, according to a different embodiment of the present invention, comprises (i) a suitable substrate to be labeled with a metal selected from the group consisting of Mn, ^{99m}Tc, ¹⁸⁶Re or ¹⁸⁸Re, (ii) a reducing agent comprising stannous ion, (iii) if desired, a stabilizing agent and/or a chelator, (iv) if desired one or more inert pharmaceutically acceptable carriers and/or formulating agents and/or adjuvants, at least one of said ingredients (i) to (iv) being
10 stored in a container having an atmosphere containing a sufficient amount of carbon monoxide, or said ingredients (i) to (iv) independently being combined, and (v) instructions for use with a prescription for reacting the ingredients of the kit with said metal in the form of a permetallate solution.

15 The preparation of the diagnostic and therapeutic pharmaceutical composition with the aid of the above mentioned kit enclosing a (biologically active) substrate can take place in two alternative embodiments. In the first embodiment the facial tricarbonyl metal compound is prepared first and then reacted with the substrate to be labeled. In the second embodiment the reduction step is carried out in the presence of the substrate to be labeled, directly leading
20 to the labeled compound.

The invention will now be described in greater detail with reference to the following specific Examples, which are offered by way of illustration and are not intended to limit the invention in any manner. Standard techniques well known in the art or the techniques specifically described below are utilized.

25

Example 1Synthesis of $[\text{Tc}(\text{CO})_3(\text{OH}_2)_3]^+$ Using SnCl_2 and Pyrophosphate or Gluceptate

Kits were prepared for performing the synthetic reactions. The first kits and results are shown in Table 1.

Table 1

Reagent/formulation	Kit A	Kit B	Kit C
Stannous chloride dihydrate ($\text{SnCl}_2 \cdot \text{H}_2\text{O}$)	0.7 mg	3.2 mg	1.6 mg
Gluceptate, sodium salt	50 mg	-	-
Pyrophosphate, sodium salt	-	11.9 mg	6 mg
pH (adjusted with 1 N NaOH)	10.5	7.3	7
Volume (0.9% saline)	1.3 mL	1.3 mL	1.1 mL
Headspace purged with CO gas	(10 cc vial)	(10 cc vial)	(10 cc vial)

Heat	100EC/30 minutes	75EC/30 minutes	75EC/30 minutes
Analysis (HPLC)	approx. 42% $^{99m}\text{Tc}(\text{CO})_3(\text{aq})_3^+$	approx. 80% $^{99m}\text{Tc}(\text{CO})_3(\text{aq})_3^+$	approx. 84% $^{99m}\text{Tc}(\text{CO})_3(\text{aq})_3^+$

Further work was performed using the following formulations. Five kits were formulated with different amounts of stannous ion and pyrophosphate ligand (potential transfer ligand for chelate attached to bioactive compounds). The five kits contained the following:

Table 2

Ingredients	Kit 1	Kit 2	Kit 3	Kit 4	Kit 5
Sodium pyrophosphate	5.0 mg	5.0 mg	10.1 mg	10.1 mg	10.3 mg
Lactose·H ₂ O	20.1 mg	20.1 mg	20.2 mg	20.1 mg	20.2 mg
SnCl ₂	0.51 mg	1.13 mg	0.50 mg	1.10 mg	3.03 mg
pH	6.1	5.0	6.5	6.0	3.4

To each lyophilized kit, 1 mL $^{99m}\text{TcO}_4^-$ in 0.9% NaCl was added and the vial (solution) was heated at 100EC for 15 minutes. Labeling was very high (>99% of the $^{99m}\text{TcO}_4^-$ was reacted with less than 1% TcO_4^- remaining, with approximately 80% of the label appearing in the desired product, the rest appearing in undesired products). The reaction products were assayed by HPLC as follows: Column: Vydac-218 TP54 C-18, 250 x 4.6 mm; Flow rate: 1 mL/minute; Mobile Phase A: 0.05 M triethylamine phosphate (TEAP), pH 2.25; Mobile Phase B: methanol; Detector: Beckman Model 170 Radiometric Detector. A gradient with linear mixing was used as shown in Table 3.

Table 3

Time Interval	Percent Mobile Phases
0-3 minutes	100% A
3-6 minutes	100% A to 75% A / 25% B
6-9 minutes	75%A / 25% B to 66% A / 34% B
9-20 minutes	66% A / 34% B to 100% B

Results are shown in Tables 4 and 5. The 3' peak is an impurity that elutes prior to the ^{99m}Tc -carbonyl. The values in Tables 4 and 5 do not add to 100% because other products, including some $^{99m}\text{TcO}_2$, were also present.

Table 4

Kit #	$^{99m}\text{TcO}_4^-$ volume	Amount of activity	Initial % Radiochemical Purity		
			3' peak	^{99m}Tc - carbonyl	$^{99m}\text{TcO}_4^-$
1	1 mL	54.5 mCi	1.9	80.5	10.6
2	1 mL	56.7 mCi	8.5	77.9	4.1
3	1 mL	48.0 mCi	1.7	79.3	10.9
4	1 mL	45.5 mCi	2.7	78.1	9.6
5	1 mL	40.0 mCi	10.3	76.7	0

Table 5

	pH at 6 Hours	mg tin detected*	6 Hour % Radiochemical Purity		
			3' peak	^{99m}Tc -carbonyl	$^{99m}\text{TcO}_4^-$
1	5.67	0.282	0	84.3	8.3
2	4.58	0.902	7.0	72.8	1.6
3	6.22	0.395	3.2	79.2	9.8
4	5.63	0.902	3.4	80.2	10.4
5	3.77	2.707	8.6	74.4	0

*The amount of tin detected after lyophilization by a method that has not been validated.

After initial labeling (reduction of $^{99m}\text{TcO}_4^-$ and formation of $\text{Tc}(\text{CO})_3(\text{OH}_2)_3^+$ complex), this compound was then reacted with an amino acid (histidine) to simulate the labeling of a chelate that may be used to attach the Tc-99m to an imaging compound such as a peptide or antibody or other bioactive molecule. For this reaction, each kit was reconstituted with 1.0 mL $^{99m}\text{TcO}_4^-$ in saline and reacted as described, then 0.2 mL of the kit was incubated with 1 mg of histidine at 75EC for one hour. Quality control was performed on the whole preparation and histidine reaction, and the pH of each reaction was measured the following morning. Results are shown in Tables 6 and 7.

Table 6

Whole Kit without Histidine						
Kit #	Amount of Activity	3' peak	^{99m}Tc -carbonyl	$^{99m}\text{TcO}_4^-$	$^{99m}\text{Tc}(\text{CO})_3\text{His}$	pH
1	73.9 mCi	6.6	84.7	5.7	-	5.60
2	72.6 mCi	14.2	61.1	0	-	4.23
3	73.0 mCi	3.1	80.7	9.7	-	6.17

4	67.0 mCi	18.3	79.3	1.6	-	5.50
5	65.5 mCi	22.9	66.5	0	-	3.65

Table 7

Histidine reaction						
Kit #	Amount of Activity	3' peak	^{99m} Tc-carbonyl	^{99m} TcO ₄ ⁻	^{99m} Tc-(CO) ₃ His	pH
1	13.6 mCi	7.0	13.0	22.4	51.6	6.74
2	13.4 mCi	67.2	0	10.0	11.6	6.44
3	13.0 mCi	8.2	0	19.0	67.9	6.82
4	11.3 mCi	33.5	0	6.1	53.8	6.50
5	10.8 mCi	86.1	0	0.1	4.3	5.94

However a mixture of what is believed to be Tc-carbonyl-ligand products was seen, where ligand was the pyrophosphate (or gluceptate if that were used instead of pyrophosphate) molecule. In the case of the Tc-CO-pyrophosphate complex mixture, addition of an excess amount histidine (3 mg) and heating for 30 minutes produced greater than 80% [Tc(CO)₃(histidine)] complex as shown by HPLC analysis. In the case where gluceptate was used in place of pyrophosphate to yield a Tc-CO-gluceptate complex, purging the solution with CO and reheating for 15 minutes increased the amount of [Tc(CO)₃(OH₂)₃]⁺. Formulation studies with the pyrophosphate kits without the addition of CO produced no such complexes.

These studies indicate that the [Tc(CO)₃(OH₂)₃]⁺ complexes can be prepared in high yields using stannous ion as the reductant. Additional transfer ligands besides pyrophosphate include phosphonates such as MDP, HEDP, HDP, etc., and ligands such as tartrate, gluceptate, etc. and carboxylic acids such as gentisic acid, ascorbic acid or oxalic acid, etc. or simple amino acids or mixed acid-amino compounds such as glycine or nicotinic acid or even histidine.

LIST OF REFERENCES

Alberto R, et al. (1994a). *J. Nucl. Biol. Med.* 38:388-90.

Alberto R, et al. (1994b). " Low CO pressure synthesis of $(\text{NEt})_2[\text{MX}_3(\text{CO})_3]$ ($\text{M} = \text{Tc}, \text{Re}$) and its Substitution Behaviour in Water and Organic Solvents." *Technetium in Chemistry and Nuclear Medicine*, No 4, Cortina International, Milano.


U.S. Pat. No. 4,232,000, Fawzi, issued Nov. 4, 1980.

U.S. Pat. No. 4,233,284, Fawzi, issued Nov. 11, 1980.

WO 98/48848

WO 96/30054 WHAT IS CLAIMED IS:


1. A method of preparing a compound of formula



wherein M is Mn, ^{99m}Tc , ^{186}Re or ^{188}Re ,

comprising reacting a metal in permetallate form with carbon monoxide and a reducing agent, wherein said reducing agent comprises stannous ion.

2. The method of claim 1 wherein said mixture further includes a stabilizing agent.
3. The method of claim 1 wherein said reducing agent forms a stannous ion.
4. The method of claim 1 wherein said reducing agent is a stannous salt.
5. The method of claim 1 wherein said reducing agent is selected from the group consisting of SnCl_2 , $\text{SnCl}_2 \cdot \text{H}_2\text{O}$, SnF_2 , SnBr_2 , $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$, SnI_2 , and SnSO_4 .
6. The method of claim 1 wherein said mixture further includes lactose.
7. The method of claim 1 wherein said mixture further includes pyrophosphate or gluceptate.
8. A method of preparing a compound of formula



wherein:

M is Mn, ^{99m}Tc , ^{186}Re or ^{188}Re ;

L_x is i) three monodentate ligands ii) one monodentate ligand and one bidentate ligand, or iii) one tridentate ligand; and

n is a charge of the ligand L_x increased with one + charge; comprising reacting a compound of formula (I) prepared according to claim 1 with ligand L_x .

9. The method of claim 8, wherein the reaction with ligand L_x takes place in the presence of a halide or a halide-like salt.
10. The method of claim 9 wherein said halide-like salt is selected from the group consisting of acetates, phosphates and sulfates.
11. The method of claim 8 wherein L_x comprises an aminopolycarboxylate.
12. The method of claim 8 wherein L_x comprises a biologically active substrate selected from the group consisting of amino acids, peptides, proteins, sugars, small receptor binding molecules and body cells.
13. The method of claim 8 wherein said method is performed between about 20EC and 150EC.
14. The method of claim 8 wherein said method is performed at about 100EC.
15. The method of claim 11 wherein said aminopolycarboxylate ligand is selected from the group consisting of diethylenetriamine-pentaacetic acid (DTPA), ethylenediaminetetraacetic acid (EDTA), 1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid (DOTA), iminodiacetic acid (IDA), nitrilotriacetic acid (NTA), and triazacyclononanetriacetate.
16. The method of claim 11 wherein said ligand is not bidentate.

17. The method of claim 11 wherein said ligand is tridentate.
18. A kit for carrying out the method of claim 1, comprising a lyophilized formulation including stannous ion, wherein said stannous ion may be in the form of a discrete molecule comprising said stannous ion plus an anion, said mixture being sealed in a container having a headspace comprising carbon monoxide.
19. The kit of claim 18 wherein said headspace is substantially pure carbon monoxide.
20. The kit of claim 18 wherein said reducing agent forms a stannous ion.
21. The kit of claim 18 wherein said reducing agent is a stannous salt.
22. The kit of claim 18 wherein said reducing agent is selected from the group consisting of SnCl_2 , $\text{SnCl}_2 \cdot \text{H}_2\text{O}$, SnF_2 , SnBr_2 , $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$, SnI_2 , and SnSO_4 .
23. The kit of claim 18 wherein said reducing agent is SnCl_2 .
24. The kit of claim 18 wherein said formulation further includes lactose.
25. The kit of claim 18 wherein said formulation further includes pyrophosphate or gluceptate.
26. The kit of claim 18 further including a metal M which is Mn, $^{99\text{m}}\text{Tc}$, ^{186}Re or ^{188}Re .
27. A kit for carrying out the method of claim 8, comprising a lyophilized formulation including stannous ion, wherein said stannous ion may be in the form of a discrete molecule comprising said stannous ion plus an anion, and a metal M which is Mn, $^{99\text{m}}\text{Tc}$, ^{186}Re or ^{188}Re .
28. The kit of claim 27 wherein said reducing agent is SnCl_2 .

29. The kit of claim 27 wherein said formulation further includes lactose.
30. The kit of claim 27 wherein said formulation further includes pyrophosphate or gluceptate.
31. The kit of claim 27 further comprising a ligand L_x which is a multidentate aminopolycarboxylate ligand.
32. The kit of claim 31 wherein L_x is not a bidentate ligand.

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
29 November 2001 (29.11.2001)

PCT

(10) International Publication Number
WO 01/089586 A3

(51) International Patent Classification⁷: **A61K 51/04**,
C07F 13/00

Elizabeth, G. [US/US]; 220 Shelburne Drive, St. Charles,
MO 63301 (US).

(21) International Application Number: PCT/US01/15670

(74) Agent: **BOONE, Jeffrey, S.**; 675 McDonnell Boulevard,
P.O. Box 5840, St. Louis, MO 63134 (US).

(22) International Filing Date: 8 May 2001 (08.05.2001)

(81) Designated States (*national*): JP, US.

(25) Filing Language: English

(84) Designated States (*regional*): European patent (AT, BE,
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,
NL, PT, SE, TR).

(26) Publication Language: English

(30) Priority Data:
09/576,960 24 May 2000 (24.05.2000) US

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

(71) Applicant (*for all designated States except US*):
MALLINCKRODT INC. [US/US]; 675 McDonnell
Boulevard, P.O. Box 5840, St. Louis, MO 63134 (US).

(88) Date of publication of the international search report:
22 August 2002

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **PIPES, David, W.**
[US/US]; 14517 Debbelham Lane, Chesterfield, MO
63017 (US). **DYSZLEWSKI, Mary, E.** [US/US]; 6 West
Walling Drive, Creve Coeur, MO 63141 (US). **WEBB,**

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/089586 A3

(54) Title: TC AND RE CARBONYL COMPLEXES FOR PERTECHNETATE AND PERRHENATE

(57) Abstract: The invention relates to novel aminocarboxylate ligands that are suitable for complexing with a radionuclide, and are useful as therapeutic agents and as imaging agents for diagnostic purposes. In accordance with the present invention, a method of preparing a compound of formula $\text{fac-[M(CO)}_3(\text{OH}_2)_3]^+$ (I) wherein M is Mn, $^{99\text{m}}\text{Tc}$, ^{186}Re or ^{188}Re , involves reacting a metal in permetallate form with carbon monoxide and stannous ion. The compound of formula (I) can be reacted with a ligand L_x , to form a compound of the formula $\text{fac-[M(CO)}_3\text{L}_x]^n$ (II) wherein M is as defined above, L_x is a monodentate or multidentate ligand or a mixture of these ligands, and n is a charge of the ligand L_x increased with one + charge. The invention also is directed to novel compounds, and kits for carrying out the disclosed methods.

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 01/15670

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 7 A61K51/04 C07F13/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07F A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, CHEM ABS Data, EMBASE, MEDLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	WO 01 25243 A (ALBERTO ROGER ARIEL ;MALLINCKRODT INC (US)) 12 April 2001 (2001-04-12) page 5, line 15-30; claims 12,15; example 3	1-32
Y	EP 0 879 606 A (SCHERRER INST PAUL) 25 November 1998 (1998-11-25)	1-32
X	abstract; claims 3-5,12,14 ----- -/--	8-17

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

4 June 2002

Date of mailing of the international search report

11/06/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax (+31-70) 340-3016

Authorized officer

Gonzalez Ramon, N

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/15670

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EGLI A ET AL: "HYDROLYSIS OF THE ORGANOMETALLIC AQUA ION FAC-TRIAQUATRICARBONYLRHENIUM(I). MECHANISM, PKA, AND FORMATION CONSTANTS OF THE POLYNUCLEAR HYDROLYSIS PRODUCTS" ORGANOMETALLICS, WASHINGTON, DC, US, vol. 9, no. 16, 1997, pages 1833-1840, XP001074762 ISSN: 0276-7333 abstract	1-7
P, Y	----- DYSZLESWKI M E ET AL: "KIT FORMULATION AND PRELIMINARY TOXICITY OF 99MTC(CO) ₃ + INTERMEDIATE: A NOVEL TECHNETIUM RADIOPHARMACEUTICAL PLATFORM" JOURNAL OF LABELLED COMPOUNDS AND RADIOPHARMACEUTICALS, SUSSEX, GB, vol. SUPPL. 1, no. 44, 2001, pages S483-S485, XP008003317 ISSN: 0362-4803 page S483	1-32
X	----- ALBERTO R ET AL: "BASIC AQUEOUS CHEMISTRY OF M(OH ₂) ₃ (CO) ₃ +(M=REMY TC) DIRECTED TOWARDS RADIOPHARMACEUTICAL APPLICATION" COORDINATION CHEMISTRY REVIEWS, ELSEVIER SCIENCE, AMSTERDAM, NL, no. 190-192, 1999, pages 901-919, XP001074720 ISSN: 0010-8545 page 905 -page 906 page 916 -page 917	8,12,13, 16,17
Y	----- SEIFERT S ET AL: "REACTIVITY OF TECHNETIUM(I) THIOETHER CARBONYL COMPLEXES TOWARDS HISTIDINE-AN EXAFS STUDY IN SOLUTION" INORGANICA CHIMICA ACTA, LAUSANNE, CH, no. 322, 2001, pages 79-86, XP001074703 ISSN: 0020-1693 abstract page 82	1-32
P, X	----- ALBERTO R ET AL: "FIRST APPLICATION OF FAC-99MTC(OH ₂) ₃ (CO) ₃ + IN BIOORGANOMETALLIC CHEMISTRY: DESIGN, STRUCTURE, AND IN VITRO AFFINITY OF A 5-HT _{1A} RECEPTOR LIGAND LABELED WITH 99MTC" JOURNAL OF THE AMERICAN CHEMICAL SOCIETY, AMERICAN CHEMICAL SOCIETY, WASHINGTON, DC, US, vol. 25, no. 121, 1999, pages 6076-6077, XP001074704 ISSN: 0002-7863 page 6077, paragraph 3; figure 1	8,12,13, 16,17
X	----- ALBERTO R ET AL: "FIRST APPLICATION OF FAC-99MTC(OH ₂) ₃ (CO) ₃ + IN BIOORGANOMETALLIC CHEMISTRY: DESIGN, STRUCTURE, AND IN VITRO AFFINITY OF A 5-HT _{1A} RECEPTOR LIGAND LABELED WITH 99MTC" JOURNAL OF THE AMERICAN CHEMICAL SOCIETY, AMERICAN CHEMICAL SOCIETY, WASHINGTON, DC, US, vol. 25, no. 121, 1999, pages 6076-6077, XP001074704 ISSN: 0002-7863 page 6077, paragraph 3; figure 1	8,13
	----- -/-	

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 01/15670

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	MEDER H-J ET AL: "METALLKOMPLEXE MIT BIOLOGISCH WICHTIGEN LIGANDEN, XLII U1 CARBONYLMETALLKOMPLEXE MIT ANIONEN VON MEHRFUNKTIONELLEN ALPHA- AMINOSAEUREN. METAL COMPLEXES WITH BIOLOGICALLY IMPORTANT LIGANDS, XLII U1 CARBONYL METAL COMPLEXES WITH ANIONS OF POLYFUNCTIONAL ALPHA-AMINO ACIDS" ZEITSCHRIFT FUR NATURFORSCHUNG, TEIL B: ANORGANISCHE CHEMIE, ORGANISCHE CHEMIE, VERLAG DER ZEITSCHRIFT FUR NATURFORSCHUNG. TUBINGEN, DE, 1986, pages 1247-1254, XP002043521 ISSN: 0932-0776 examples 2A,2B	8,13
X	BAMFORD C H ET AL: "EVIDENCE FOR THE FORMATION OF THE TRIAQUATRICARBONYLMANGANESE(L) CATION AND RELATED DERIVATIVES FROM PENTACARBONYLCHLOROMANGANESE" JOURNAL OF THE CHEMICAL SOCIETY, DALTON TRANSACTIONS, CHEMICAL SOCIETY. LETCHWORTH, GB, no. 1, 1978, pages 4-8, XP002043519 ISSN: 1472-7773 abstract	8,13
P,Y	WO 01 00637 A (ALBERTO ROGER ;DYSZLEWSKI MARY M (US); BUGAJ JOSEPH E (US); MALLIN) 4 January 2001 (2001-01-04) abstract; claims 1,4,6,8,9,15; examples 2,3	1-32

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/15670

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0125243	A	12-04-2001	AU 1134401 A WO 0125243 A1	10-05-2001 12-04-2001
EP 0879606	A	25-11-1998	EP 0879606 A1 AT 211924 T AU 7141398 A BR 9809409 A DE 69803205 D1 DK 1019095 T3 EP 1019095 A1 HU 0003255 A2 JP 2002512616 T NO 995160 A NZ 337303 A PL 336382 A1 WO 9848848 A1 US 6344178 B1	25-11-1998 15-02-2002 24-11-1998 13-06-2000 21-02-2002 06-05-2002 19-07-2000 28-02-2001 23-04-2002 13-12-1999 22-12-2000 19-06-2000 05-11-1998 05-02-2002
WO 0100637	A	04-01-2001	AU 6488900 A EP 1192166 A1 WO 0100637 A1	31-01-2001 03-04-2002 04-01-2001

This Page Blank (uspto)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☒ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

This Page Blank (uspto)